

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A23K 1/16, 1/18</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/65327</b> <b>(43) International Publication Date:</b> 23 December 1999 (23.12.99)
<b>(21) International Application Number:</b> PCT/EP99/04224 <b>(22) International Filing Date:</b> 17 June 1999 (17.06.99)  <b>(30) Priority Data:</b> 98304802.6      17 June 1998 (17.06.98)      EP  <b>(71) Applicant (for all designated States except US):</b> GIST-BROCADES B.V. [NL/NL]; Wateringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BEUDEKER, Robert, Franciscus [NL/NL]; Boomkwekerij 31, NL-2635 KC Den Hoom (NL). COUTTEAU, Peter [BE/BE]; Inve Technolo- gies N.V., Oeverstraat 7, B-9200 Baasrode (BE).  <b>(74) Agents:</b> WRIGHT, Simon, Mark et al.; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5LX (GB).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
<b>(54) Title:</b> MICROBIAL ARACHIDONIC ACID (ARA) FOR USE IN MARINE FEED  <b>(57) Abstract</b>  A marine feed composition is described comprising microbially derived arachidonic acid (ARA) or ARA in the form of a triglyceride. The ARA is suitably produced by a fungus, such as of the genus <i>Mortierella</i> , although the feed composition itself is free of microbial cells. These forms of ARA have been found to give better growth and pigmentation promotion in marine organisms (shrimps and fish) than corresponding phospholipid forms of ARA from fish oil. The ARA can be in the form of an oil, e.g. an oil-in-water emulsion or may first be fed to larvae, rotifers or nauplii which are themselves included in a composition as "live" feed for larger organisms.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

MICROBIAL ARACHIDONIC ACID (ARA)  
FOR USE IN MARINE FEED

Field of the invention

The present invention relates to the use of microbially derived arachidonic acid  
5 (ARA) for use in marine feed compositions, for feeding aquatic (e.g. marine)  
animals and for use in aquaculture. The ARA can be derived from a single cell  
organism, and thus may be in the form of a triglyceride.

Background of the invention

Arachidonic acid (ARA) is one of a number of Long Chain Poly-Unsaturated  
10 Fatty Acids (LC-PUFA's). Chemically it is *cis*-5,8,11,14 eicosatetraenoic acid (20:4)  
and belongs to the (n-6) family of LC-PUFA's.

Arachidonic acid is a major precursor (in fish as well as in higher terrestrial  
vertebrates) of a wide variety of biologically active compounds, known collectively  
as eicosanoids, a group comprising prostaglandins, thromboxanes and leukotrienes.  
15 Marine fish, shrimp and various terrestrial animals, such as cats, lack  $\Delta$ -5 desaturase  
activity and so are not able to convert the fatty acid 18:2(n-6) (linoleic acid) to  
ARA. Since arachidonic acid is essential to produce eicosanoids, it is an essential  
fatty acid for these animals and so must be provided in their diets.

The ratio of eicosapentaenoic acid (EPA) {20:5(n-3)} to docosahexaenoic acid  
20 (DHA) {22:6(n-3)} to arachidonic acid {20:4(n-6)} in the diets of these animals can  
be important since a relative excess of EPA can prevent the production of  
eicosanoids from arachidonic acid.

-2-

Normal diets for marine fish and shrimp contain variable amounts of arachidonic acid of fish oil origin. The content of arachidonic acid in known fish oils is variable but usually relatively low, about 1% of total fatty acids present (Sargent, J.R., McEvoy, L.A. and Bell, J.G.: Aquaculture 155 (1997) p.117-127).

- 5 The ability to grow marine fish larvae is often determined to a large extent by the nutritional composition of the diet. It is difficult to obtain desirable ratios of eicosapentaenoic acid, docosahexaenoic acid and arachidonic acid using available raw materials such as fish oil, animal offal and the like since the LC-PUFA levels are highly variable even similar sources and a substantial part of arachidonic acid  
10 present is destroyed in the process of sterilising these raw materials.

- One of the aims of the invention is to provide a source of arachidonic acid, of reproducible quality, to enable the aquaculture feed industry to find a satisfactory level of arachidonic acid in the diet (e.g. live feeds) for fish larvae and shrimp as well as the right ratio of arachidonic acid, eicosapentaenoic acid and  
15 docosahexanoic acid for these animals. Nearly all marine culture (mariculture) production systems rely on live feeds, principally the rotifer *Brachionus plicatilis* and nauplii of the brine shrimp *Artemia*, although alternative live feeds do exist.

- Enrichment of live feed with eicosapentaenoic acid and docosahexaenoic acid from fish oil emulsions have been described (Sargent *et al, supra*). They also describe the  
20 possibility of using freeze-dried or spray-dried cells of algae and fungi containing high amounts of docosahexaenoic acid for the enrichment of live feeds.

- Sargent *et al* also suggests arachidonic acid, but in the form of a phospholipid. Phospholipids of arachidonic acid are a natural source for flatfish and shrimp in their live feeds as this is the form that is present in fish oil. The phospholipid form  
25 of arachidonic acid is less susceptible to oxidation (in comparison with free arachidonic acid) and so is preferred by the aquaculture industry.

-3-

WO-A-90/07283 (Suntory) refers to feeds for fish and shellfish containing fungal cells which contain n-3 fatty acids (e.g. EPA, DHA) or esters thereof. However, again these sources are often low in ARA and furthermore the ARA is often present as a phospholipid.

- 5 It has been found however that by using microbially derived ARA a much higher content of ARA is achievable inside the marine animal. A marked increase in enrichment in ARA has been found in marine animals fed with feeds that contain ARA derived from single cell organisms.

#### Description of the Invention

- 10 Thus, according to a first aspect of the present invention there is provided a marine feed composition comprising one or more (marine) feed component(s) and/or ingredient(s) and microbially derived arachidonic acid (ARA). The ARA can thus be derived from a single cell organism, such as a microorganism, which it will be appreciated excludes fish oil (or marine) sources. Microbially derived ARA  
15 is usually in triglyceride form and so the invention also relates to a marine feed composition comprising ARA in the form of a triglyceride.

- Preferably the feed will be (substantially) free of ARA in the form of phospholipids. The invention thus may avoid the use of phospholipid ARA, which is the form that ARA occurs in marine sources (e.g. fish oil). To thus feed  
20 marine animals with a form of ARA different from the natural source, in other words a form that would not normally be ingested by the organism, seems counter-intuitive. Nevertheless, the Applicant has found that this can lead to unexpected benefits.

- The ARA will usually be present as or in an oil. Thus, this oil may be derived  
25 from or produced by a single cell organism, such as from a microbial source. The

-4-

oil may have at least 10%, such as at least 20%, preferably at least 30%, advantageously at least 35% and optimally at least 40% ARA (based on the total weight of fatty acids). The oil may have an ARA content of at least 250, preferably at least 300, optimally at least 350 g/kg oil. The oil suitably has a  
5 minimum triglyceride content of 90%, eg. at least 95%, optimally at least 97%. Of the ARA present, preferably at least 85%, such as at least 90%, and optimally at least 95% is in the form of triglycerides.

Preferably the ARA is not still inside cells, for example the microbial cells that had produced the ARA. Thus, the marine feed of the invention can be (substantially)  
10 free of single cell organisms or microbes such as from which the ARA is derived or produced by. This can not only make the feed composition easier to formulate but, surprisingly, the ARA is found to be more stable once extracted from the cells that produced the ARA.

Although ARA has been suggested for use in marine feeds, the tendency has been  
15 to use ARA in a phospholipid form, usually from a marine source. This is not only because the phospholipid form is the naturally occurring form for marine animals but also since it is less susceptible to oxidation. However, despite this prejudice, the applicant has found that microbially derived ARA (or ARA in the form of a triglyceride) can provide advantages over phospholipid forms of ARA.  
20 In particular there can be a higher uptake and use of the ARA by the marine organism to which the feed composition of the invention is fed. Furthermore, with high quality microbial ARA the oxidation problem can be overcome or at least mitigated.

A further advantage of microbial ARA is that the oil can have a higher ARA  
25 content than, say, fish oil, which means less microbial oil can be used to deliver the same quantity of ARA (resulting in cheaper transport, storage and feed costs).

-5-

The microbial (or single cell) source from which the ARA is derived or produced by is suitably a microorganism, for example one in a filamentous form, such as a fungus or an algae. Preferred fungi are of the order *Mucorales*. For example, the fungus may be of the genus *Mortierella*, *Phycomyces*, *Blakeslea* or *Aspergillus*.

- 5 Preferred fungi are of the species *Mortierella alpina*, *Blakeslea trispora* and *Aspergillus terreus*.

Preferred algae are dinoflagellates and/or belong to the genus *Crypthecodinium*.

Preferred algae are of the species *Crypthecodinium cohnii*.

- 10 These ARA sources produce predictable quantities of ARA, unlike marine sources whose PUFA content can vary according to species and the processing techniques used.

- 15 The ARA is preferably derived from or produced by the microbial (or single cell) source by fermentation. The microbes may be grown in (large scale) fermentors. Fermentation can occur when the microorganisms are present in an aqueous composition comprising an assimilable carbon source (for example, glucose or maltodextrin) and a suitable nitrogen source (for example, ammonium ions, such as from ammonium sulphate or yeast extract).

- 20 During or after fermentation one obtains a fermentation "broth" which comprises the microorganisms and a medium, e.g. an aqueous solution. These can then be separated (eg. by filtration, centrifugation or other solid-liquid separation technique) to give the aqueous solution (eg. in the form of a filtrate) and the microorganisms (referred to as a biomass). The ARA (and other PUFAs, if necessary) can then be extracted and/or isolated from the biomass. This can be achieved using a solvent, for example, an apolar solvent or any other suitable solvent that is capable of dissolving ARA. In particular, this can be a C<sub>3-8</sub> alkane, such as hexane.
- 25

-6-

Before the ARA is extracted, the biomass can be subjected to mechanical, chemical and/or physical processing. For example, the biomass may be heated and/or pasteurised. The biomass may be processed into a granulate form, for example by extrusion.

- 5 The ARA, as explained earlier, is preferably present in the marine feed as an oil. This oil may be present as an emulsion. Thus, the ARA may be provided as an oil in a oil-in-water emulsion. The emulsion preferably contains an emulsifier, and may additionally contain an antioxidant. Lecithin may fulfill both functions.

- 10 The compositions of the invention may be used as feeds for aquatic animals and, generally, in aquaculture. Suitable marine organisms to which the composition can be fed are salt or seawater organisms. Suitably the organisms are coldwater organisms.

- 15 The marine organism to be fed the composition may be larvae (e.g. of the subPhylum *Crustacea*), rotifers (such as of the Phylum *Rotifera*) or nauplii (e.g. larvae of *Artemia*). Preferred rotifers are of the species *Brachonius plicatilis*. All these organisms are relatively small and so may themselves be present in the marine feed for digestion by larger organisms. In this case these organisms are one of the feed components or ingredients and so constitute a "live" feed. In this case the ARA may be already inside these organisms or it may be present elsewhere in  
20 the composition.

Other suitable marine organisms that may be fed the compositions of the invention include the copepods (for example of the class *Copepoda*), for example of the class *Tisbe*, *Acartia*, *Artemia* and/or *Eurytemora*.

- 25 Marine organisms specifically contemplated belong to the Phylum *Mollusca*, for example of the class *Bivalvia*, subclass *Protobranchia* and *Lamellibranchia* (which



-7-

includes *Mytilus* and *Ostrea*, mussels and oysters). Also included are organisms of the class *Cephalopoda*, such as subclass *Coleoida*, for example *Decapoda*, *Sepia*, *Loligo* (including squids and cuttlefish) and *Octopoda* (octopus).

- Other organisms are of the subPhylum *Crustacea*, such as division *Astacus*, which  
 5 includes crayfish, and order *Amphipoda*, such as division *Gammarus*, which includes prawns and shrimps.

- The marine organisms can therefore include non-vertebrates such as octopus and squid, prawns (*Leander*) and shrimps (*Crangon*). Also included are fish, or other vertebrate organisms, such as herring, turbot (e.g. *Scophthalmus maximus*), halibut,  
 10 salmon (Atlantic, Chinook, Coho, Pacific), trout (such as Rainbow trout, *Oncorhynchus mykiss*), bream (such as sea bream, *Sparus aurata*), sole (such as common sole, *Solea solea* L, Senegal sole, *Solea senegalensis* Kaus), bass (such as sea bass, *Dicentrarchus labra*), flounder (*Parlichtrys*, such as Japanese flounder, *Parlichtrys divaceus*), Yellow tail, Milkfish and Arctic Charr.

- 15 Also included are shellfish such as lobster, mussels, oysters, clams, prawns (*Macrobrachium*, such as *rosenbergii*), shrimps (*Penaeus*, such as *japonicus*, Kurum shrimp and *vannamei*, white leg shrimp, *monodon* and *setiferus*).

Preferred shrimps are those of the class *Artemia*. In particular, nauplii of *Artemia* can be fed or included in the compositions of the invention as a "live" feed.

- 20 Live feeds, where an edible marine organism is provided in the food for consumption by a larger organism, are particularly suitable for young organisms or those at an early stage of development, e.g. larvae, for example up to 3 days, such as up to 5 days or even up to 10 days old. This is because such organisms may not be able to assimilate prepared diets effectively. The inclusion of live organisms  
 25 in the feed will usually be in addition to various or standard feed components

-8-

and/or ingredients. Live nauplii, e.g. of brine shrimp or *Artemia* are particularly suitable for larvae.

Other components or ingredients may include, for example, fish meal, soya meal, wheat meal binders and/or stabilizers, and if necessary minerals and/or vitamins.

- 5 The feed may be in the form of pellets and/or granules. Larvae (e.g. of fish and crustaceans) may initially be fed on live feeds, and later on in development non-live feeds.

To give some guidance on the amounts of various materials that can be included, these may be as follows:

- |    |  |                                    |
|----|--|------------------------------------|
| 10 | fish meal                                      | 30 to 60%, e.g. 40 to 50%;         |
|    | enzymatically digested fish meal (concentrate) | 1 to 15%, e.g. 5 to 10%;           |
|    | soya and/or wheat meal                         | 5 to 25%, e.g. 10 to 18% (each) ;  |
|    | minerals and/or vitamins                       | 0.5 to 4%, such as 1 to 2% (each). |

- For many marine organisms, in particular fish, the diet can contain 10 to 40% lipids. It may also contain from 35 to 50% protein. Suitably the amount of phosphorus available to the organism is less than 0.8%. This diet may contain natural or artificial carotenoids, vitamins and/or minerals.
- 15

- It is preferred that the amount of ARA present in a (e.g. non-live) feed is from 0.1 to 5%, such as from 0.1 to 2%. Suitably the total n-6 PUFA content can be from 1 to 15% of the diet (by weight).
- 20

For a live feed the ARA content may be from 5 to 30%, for example 10 to 25%, optimally from 15 to 20%. The total content of n-6 PUFAs may then be from 5 to 50%, e.g. 5 to 30%. Here the ARA may be inside (having already been consumed by) the live edible marine organisms.

-9-

If necessary, the marine feed may contain other polyunsaturated fatty acids (PUFAs), for example a C18, C20 and/or C22  $\omega$ 3 or  $\omega$ -6 PUFA. Other PUFAs contemplated include for example docosahexaenoic acid (DHA) (from algae or fungi, such as the algae *Cryptothecodinium* or the fungus *Thraustochytrium*),  
5 gamma-linolenic acid (GLA), and/or EPA. Any of these PUFAs may be provided individually or may be present in the oil in which the ARA is contained.

If EPA is provided in addition to ARA, then the ratio of EPA:ARA can be from 1:0.5 to 1:2, such as from 1:1 to 1:1.6. If DHA is present, then the DHA:ARA ratio may be from 1:0.2 to 1:1, such as from 1:0.25 to 1:0.7. When both EPA and  
10 DHA are present then the DHA:EPA:ARA ratio may vary from 3-5:0.5-2.0:1.0-2.0 (e.g. if the feed is not "live") or from 1.0-2.0:0.7-1.3:0.7-1.3 for a "live" feed.

A second aspect of the present invention relates to a process for the preparation of a marine feed composition, the process comprising mixing microbially derived ARA, or ARA in the form of a triglyceride, with one or more marine feed  
15 component(s) and/or ingredient(s). These may include fish meal and materials of vegetable origin and have been described in detail earlier.

The feed compositions may, for example, be "live" compositions containing live multicellular marine organisms, for example larvae, nauplii or rotifers as  
20 previously described.

This aspect of the invention therefore includes the supplementation of an existing marine feed composition with the ARA.

A third aspect of the present invention relates to the use of microbially derived ARA, or ARA in the form of a triglyceride (or marine feed compositions  
25 comprising either of these forms of ARA) for use in promoting the growth and/or pigmentation in a marine organism. This use may also extend to the

-10-

supplementation of a marine diet, for example as a nutritional supplement. The use also contemplates the assistance of the marine organism in producing eicosanoids.

5 A fourth aspect of the present invention relates to a process of feeding a marine organism with a marine feed composition of the first aspect.

A fifth aspect of the invention relates to a process for promoting growth and/or pigmentation in a marine organism, the process comprising feeding the organism with a marine feed composition of the first aspect.

10 Preferred features and characteristics of one aspect of the present invention are suitable for another aspect *mutatis mutandis*.

The invention will now be described, by way of example, with reference to the following Examples, which are provided for illustration, and are not to be construed as being limiting.

COMPARATIVE EXAMPLES 1 and 2 and EXAMPLE 3Enrichment of *Artemia* feed with microbial arachidonic acid (ARA)

Crude arachidonic acid (ARA) oil was produced by the fungus *Mortierella alpina* using the protocol of Example 1 of the International Application No.

- 5 PCT/EP97/01446 (WO-A-97/36996). This oil is available from Gist-brocades B.V. under the trade mark VEVODAR™. The oil was then ready for formulation of arachidonic acid rich emulsions using lecithin (to give a final concentration of lecithin of 10%).

- 10 An emulsion of the ARA oil was formulated to give a ratio of docosahexaenoic acid (DHA), eicosapentaenoic acid (EPA) and arachidonic acid (ARA) of 4/1/1.5 (Example 3) while another emulsion (Comparative Example 2) was formulated to a DHA:EPA:ARA ratio of 2.5/1/0.2 by using fish oil instead of the microbial ARA oil. The two emulsions were then tested for stability by centrifuging the emulsions for 15 minutes at 3000 rpm. Both emulsions were stable in this
- 15 centrifuge test, meaning that no separation into different layers was observed.

*Artemia* cysts were hatched in seawater at a temperature of 28°C. After 24 hours from hatching, the nauplii were harvested and transferred to an enrichment tank at a density of 250 nauplii/ml.

- The two emulsions were then used (each in duplicate) to supplement a standard nauplii diet:
- 20

- a) enrichment with 300ppm microbial arachidonic acid (ARA) emulsion (Example 3); and
- b) enrichment with 300 ppm fish oil emulsion (Comparative Example 2).

After subsequent enrichment, samples of the nauplii were taken to perform fatty acid analysis using the FAME (fatty acid methyl esters) methodology as described by Lepage and Roy, J.Lip.Res. 25:1391-6 (1984) to determine their PUFA content. As a control some nauplii were grown using a standard feed without either  
 5 emulsion (no ARA, Comparative Example 1).

The levels of docosahexaenoic acid, eicosapentaenoic acid and arachidonic acid in enriched nauplii and control nauplii are shown in Table 1.

Table 1: Levels of docosahexaenoic acid, eicosapentaenoic acid, arachidonic acid and the sum of all (n-3) and (n-6) fatty acids in *Artemia* nauplii (after enrichment).

10	Enrichment Protocol (Example)	Docosahexaenoic acid (DHA, n-3) content (mg/g dry weight)	Eicopentaenoic acid (EPA, n-3) content (mg/g dry weight)	Arachidonic acid (ARA,n-6) content (mg/g dry weight)	Sum (n-3) + (n-6) fatty acids (mg/g dry weight)
	1. Control (none)	0.1	7.9	1.6	20.7
15	2. Fish oil emulsion	28.3	23.5	4.6	84.6
	3. Microbial ARA emulsion	24	15.6	16.1	83.3

20 The results show that arachidonic acid oil made by *Mortierella alpina* can be used to enrich *Artemia* feed. This resulted in an altered ratio of docosahexaenoic acid, eicosapentaenoic acid and arachidonic acid in the fed nauplii. The ratio of fatty acids provided by the microbial oil was more suitable to increasing growth and pigmentation in aquatic species by significantly increasing the ARA content in the  
 25 organisms. These ARA-enriched nauplii can be used themselves as live feed ingredients in feed compositions for larger marine animals.

The same procedure was repeated except using rotifers (*Brachionus plicatilis*)

instead of *Artemia* nauplii.

#### COMPARATIVE EXAMPLES 4 and 5 and EXAMPLE 6

##### Effect of ARA enriched diet on growth parameters of tiger shrimp

- Four day old *Penaeus monodon* (giant tiger shrimp) postlarvae were obtained from a commercial source and were grown in tanks containing disinfected seawater at a density of 20 postlarvae/l at a temperature of 28°C. ARA enriched *Artemia* nauplii and control nauplii (from Examples 2 and 3) were fed ad libitum to the tanks 4 times a day (as described by Rees, J.F., Cure, K., Piyatiratitivorakul, Sorgeloos, P and Menasveta, P in Aquaculture 122 (1994) 193-207) for 10 days.
- The increase in total mass of the larvae was taken as the indicator for growth. The dry weight (of total biomass) was measured at the beginning and the end of the experiment and the increase in weight is shown in Table 2.

- Table 2: Growth of monodon postlarvae fed *Artemia* nauplii enriched with ARA from a microbial source in comparison with non-enriched nauplii and nauplii enriched with fish oil (nauplii prepared according to Examples 2 and 3).

Example	Enrichment protocol	Increase in Shrimp dry weight (mg)
4	Control (none)	30
5	Fish oil emulsion (Ex. 2)	92
6	Microbial ARA emulsion (Ex. 3)	135

- The results clearly show the favourable effects of microbial ARA enriched *Artemia* nauplii on the growth of the larvae.

At the end of the experiment the then 15 day-old postlarvae were subsequently

-14-

lyophilized. The fatty acid composition of the larvae was analysed using the standard technique described in the previous Example. Results are expressed as mg FAME (fatty acids methyl esters)/g dry weight of shrimp.

5 Table 3: Fatty acid composition of tiger shrimp postlarvae fed *Artemia* enriched with microbial ARA in comparison with nauplii enriched with a fish oil emulsion and a negative control (i.e. nauplii not enriched with PUFA's).

Fatty acid	Comparative Example 4 Negative control	Comparative Example 5 Fish oil	Example 6 microbial-ARA oil
ARA	2.8	3.4	9.7
DHA	0.4	8.5	8.3
10 EPA	3.1	9.7	9.5

The results demonstrate the marked increase in ARA content in postlarvae using a mycrobial source of ARA.

#### COMPARATIVE EXAMPLES 7 and 8 and EXAMPLE 9

##### Effect of ARA enriched diet on growth of white bass larvae

- 15 A similar experiment to that described in Examples 4 to 6 was repeated except using larvae of white bass. The methodology used was described in detail by J.A. Clawson and T.T. Lovell in *Aquaculture* 108 (1992): 125-134. The results obtained show that feeding white bass larvae with *Artemia* nauplii enriched in microbial ARA improved growth in comparison with control *Artemia* nauplii enriched with
- 20 fish oil.



CLAIMS

1. A marine feed composition comprising one or more feed component(s) and/or ingredient(s) and microbially derived arachidonic acid (ARA).
- 5 2. A marine feed composition comprising one or more feed component(s) and/or ingredient(s) and ARA where the ARA is in the form of a triglyceride and/or the composition is substantially free of ARA in phospholipid form.
3. A composition according to any preceding claim which is  
10 substantially free of microorganisms that produced, or are capable of producing, ARA.
4. A composition according to any preceding claim wherein the ARA is produced by or derived from a fungus or an algae.
5. A composition according to claim 4 wherein the fungus is of the  
15 order *Mucorales*.
6. A composition according to claim 5 wherein the fungus is of the genus *Mortierella*, *Phycomyces*, *Blakeslea* or *Aspergillus*.
7. A composition according to any of claims 4 to 6 wherein the fungus is the species *Mortierella alpina*.
- 20 8. A composition according to any preceding claim wherein the ARA is present as an oil.
9. A composition according to claim 8 wherein the ARA content in the oil is at least 20%.
10. A composition according to claim 8 or 9 wherein the oil comprising  
25 ARA is formulated as an oil-in-water emulsion.
11. A composition according to any preceding claim which is a "live" feed composition and comprises edible marine organisms.
12. A composition according to claim 11 wherein either the edible marine organisms that are present as a live feed, or the marine organisms to which

-16-

the composition is fed, are larvae, rotifers or nauplii.

13. A composition according to claim 12 wherein the organisms are of the subPhylum *Crustacea* or Phylum *Rotifera*.

14. A composition according to claim 12 or 13 wherein the ARA is  
5 present in the edible marine organisms in the live feed.

15. A composition according to any preceding claims which additionally comprises a C<sub>18</sub>, C<sub>20</sub> or C<sub>22</sub>  $\omega$ -3 or  $\omega$ -6 polyunsaturated fatty acid (PUFA).

16. A composition according to claim 15 wherein the PUFA comprises  
10 DHA, GLA and/or EPA.

17. A process for preparing a marine feed composition, the process comprising mixing either microbially derived arachidonic acid (ARA) or ARA in the form of a triglyceride with one or more feed component(s) and/or ingredient(s).

18. The use of microbially derived ARA, ARA in the form of a triglyceride, or a marine feed composition according to any preceding claim for promoting growth and/or pigmentation in a marine organism.

19. A process for promoting growth and/or pigmentation of a marine organism, the process comprising feeding the organism a marine feed composition  
20 as defined in any of claims 1 to 16.

20. A process of feeding a marine organism, the process comprising supplying the organism with a feed composition according to any of claims 1 to 16.

21. A process according to claim 19 to 20 wherein the marine organism  
25 is of the class *Copepoda* or *Cephalopoda*, Phylum *Mollusca*, subPhylum *Crustacea*, *Leander* or *Crangon*.

22. A process according to any of claims 19 to 21 wherein the organism is a mussel, oyster, squid, cuttlefish, octopus, crayfish, prawn, shrimp, herring, turbot, halibut, salmon, trout, bream, sole, bass, flounder, charr, lobster or clam.